



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

made by Dr. W. R. Miller, surgeon to the West Riding convict prison. Dr. Miller experimented on about four thousand prisoners for thirteen years, and obtained results that differ sensibly from those of Mr. Hansen; for he found that the season of maximum increase in weight in adults is from April to August, and the period of minimum increase in adults from September to March. Dr. Miller found the body became heavier in summer, and lighter in winter; and he attributes the loss of weight to the more active excretion of carbonic-acid gas in the colder months.

DIFFERENT PHYSIOLOGICAL SENSES FOR HEAT AND COLD.

IN connection with the researches of Professor Hall and Dr. Donaldson of Johns Hopkins university, recently given in *Mind*, it will be of interest to state that Mr. A. Herzen has lately published in the *Archiv für physiologie*¹ the results of a series of experiments showing that the physiological sense of cold is different from that of heat. His attention was first directed to the subject by a simple incident, the verification of which may be readily made by any observer. Awakening one night, he found one of his arms lying without the bed-clothes, 'alseep;' in touching it with the other hand, he perceived a distinct sensibility to warmth, while that of touch was gone. Bringing his arm, however, in contact with cold substances, he was surprised to find no sensation.

Pursuing the subject further, he produced artificially this condition of semi-paralysis by the compression of nerve-trunks, and by experimentation discovered that the sensibility to cold remained a short time after tactile impressions had disappeared, and that the sensibility to warmth remained much longer, but not quite as long as the power of detecting pain; also that the impressions of warmth require more time for transmission to the brain than those of cold, bearing, in fact, the same relations to each other as the sense of pain does to that of cold. These results were further supplemented by observations on a person with complete and permanent tactile anaesthesia of the legs, but in whom the sense of pain remained normal. The subject was able to distinguish quite well the differences in temperature between 150° F. and 81° F., which was the normal temperature of the surface of the leg. Below the latter temperature, however, no sensation was produced, not even by the contact of ice on the inner side of the thigh. Other cases showed the same peculiarities, in which,

with the disappearance of tactile sensibility, the susceptibility to cold was also lost, while that to warmth yet remained.

Vivisection experiments upon cats and dogs lead the author to the following conclusions: 1. The so-called sense of heat and cold is composed in reality of two senses quite independent both anatomically and physiologically; 2. Observations on healthy and diseased subjects show that the sensations of heat and cold are transmitted through different nerves, by different routes, and to different brain-centres; 3. The gyrus sigmoides contains the centre (or the centripetal branches leading thereto) of touch and cold perceptions; 4. These sense-perceptions are transmitted through the posterior columns of the spinal cord, while those of the senses of pain and warmth are conveyed through the gray substance.

Although the senses of cold and touch on the one hand, and heat and pain on the other, seem to be more nearly related, yet one cannot unite them, or consider the different perceptibilities of heat and cold mere modifications of those of touch and heat. The researches of Blix and others have demonstrated the existence of separated, isolated, irregularly distributed points upon the body, of which one may be only sensible to cold, another to warmth, and a third to touch. Doubtless most persons have noticed the different degrees of susceptibility of different parts of the body to heat and cold: the author points out striking examples of such.

RAINFALL IN SOUTH AFRICA.¹

LITTLE has been known until recently on the subject of rainfall in South Africa, taken broadly over the whole country, although observers have for many years been keeping records at isolated stations. There has been for many years a meteorological commission in existence at Cape Town; and in the report for 1883 an interesting table was published, giving the means, monthly and yearly, at all stations where records have been kept for at least five years, with the altitude above sea-level, and the latitude and longitude for each station.

From these data Mr. Tripp has prepared a map of South Africa, with the idea of showing the distribution of the total yearly rainfall. The curves divide the area into districts, where the mean yearly rainfall is—

(1) Under	5 inches.	
(2) From	5 "	to 10 inches.
(3) " "	10 "	" 20 "
(4) " "	20 "	" 30 "
(5) Above	30 "	

There are doubtless, particularly along the moun-

¹ *Ueber die spaltung des temperatursinnes in zwei gesonderte sinne*, xxxviii. p. 93, December, 1885.

¹ Abstract of an article by William B. Tripp, in *Symons's meteorological magazine*.